

Quantum Origins of Molecular Recognition and Olfaction in *Drosophila*

Eric R. Bittner,¹ Adrian Madalan,² Arkadiusz Czader,³ and Gregg Roman²

¹⁾Depts. of Chemistry and Physics, University of Houston, Houston TX, 77204

²⁾Dept. of Biology and Biochemistry, University of Houston, Houston TX, 77204

³⁾Dept. of Chemistry, University of Houston, Houston TX, 77204

The standard model for molecular recognition of an odorant is that receptor sites discriminate by molecular geometry as evidenced that two chiral molecules may smell very differently. However, recent studies of isotopically labeled olfactants indicate that there may be a molecular vibration-sensing component to olfactory reception, specifically in the spectral region around 2300 cm^{-1} . Here we present a donor-bridge-acceptor model for olfaction which attempts to explain this effect. Our model, based upon accurate quantum chemical calculations of the olfactant (bridge) in its neutral and ionized states, posits that internal modes of the olfactant are excited impulsively during hole transfer from a donor to acceptor site on the receptor, specifically those modes that are resonant with the tunneling gap. By projecting the impulsive force onto the internal modes, we can determine which modes are excited at a given value of the donor-acceptor tunneling gap. Only those modes resonant with the tunneling gap and are impulsively excited will give a significant contribution to the inelastic transfer rate. Using acetophenone as a test case, our model and experiments on *D. melanogaster* suggest that isotopomers of a given olfactant give rise to different odorant qualities. These results support the notion that inelastic scattering effects play a role in discriminating between isotopomers, but that this is not a general spectroscopic effect.

I. INTRODUCTION

The general model for detection is that the response is triggered by the transfer of an electron from a donor (D) to an acceptor (A) within the receptor site by the presence of an olfactant molecule that provides a bridge between the two. In the absence of the odorant, the distance between D and A is too great and electron transfer is inefficient. Placing an olfactant (B) between the two allows the electron transfer to occur either as a single coherent scattering event from D to A, or as a sequence of two incoherent hops, first from D to B then from B to A. This simple model is consistent with the standard “swipe-card” model for odor detection since implicit in this is that B must fit into some sort of pocket and be in a correct alignment between D and A in order for the charge transfer process to occur.

An interesting extension to this paradigm is that molecular shape may not be the sole deciding factor in scent recognition. Indeed, it has been observed recently that fruit flies (*D. melanogaster*) can naïvely discriminate between several isotopomers that have deuteriums substituted for hydrogen¹. These flies can also be trained to associate a specific odorant with an electric shock resulting in the specific avoidance of the conditioned odorant². Flies trained in this manner can discriminate deuterated from hydrogenated isotopomers, indicating an ability to perceive differences in these odorants. Mutant flies that lack the orco olfactory co-receptor gene are broadly anosmic and fail to discriminate between isotopomers, indicating the behavioral discrimination of isotopomers relies on olfactory perception. Lastly, using the negatively-reinforced learning paradigm, *Drosophila* has been shown capable of generalizing a trained odorant to another that shares a similar vibrational spectrum. For example, when deuterated d5-benzaldehyde is paired with electric shocks, the flies will subsequently

avoid fully-deuterated d17-octanol (vs. undeuterated h-octanol), suggesting that the C-D vibrational mode is a salient feature of odorant perception in this species.

A controversial and speculative explanation of these observations by Turin is that there is an spectroscopic component to olfaction.^{3,4} Turin’s model supposes that the bridge facilitates inelastic electron scattering and as such the olfactory response can be predicted by comparing infra-red (IR) spectra of various olfactants. Isotopic substitution of H for D shifts the CH stretching frequency from the $2850 - 3100\text{ cm}^{-1}$ range into the 2300 cm^{-1} range. Few molecules absorb in this IR region and there is little or no biological need or evolutionary pressure that we know for detecting deuterated compounds. Needless to say, this theory has been met with considerable skepticism since it runs contrary to more commonly held model in which the geometric shape and chemical nature of the olfactant are the primary components of olfactory reception.⁵ Moreover, many of the claims in Ref. 3 and Ref. 4 were shown to be inconsistent with psychophysical tests performed on human subjects by Keller and Vosshall.⁶ For example, in this study human subjects could not discriminate by smell between deuterated and non-deuterated forms of acetophenone.

While the olfactory receptors and the co-receptor have been identified and is believed to be a multi-helix transmembrane protein with the binding sites on the periplasmic domain and a G protein-binding site on the cytoplasmic domain, information concerning the molecular structure at the olfactant binding site remain illusive, but appear to involve odorants binding to Cu(II) and Zn(II) ions bound to the protein loop.⁷ We can assume that at the heart of the process, a charge is transferred from a donor to an acceptor within the receptor site itself. In the absence of an odorant, the distance between the donor and acceptor is too great to allow efficient electron transfer. When the odorant is present, it acts like a




FIG. 1. Calculated IR Spectrum (a) and projection of the energy gradients (in atomic units) (b) of ACP⁺ and its deuterated isotopomers. Color key: —: hACP, —: d3ACP, —: d5ACP, —: d8ACP.

bridge between the donor and acceptor allowing for more efficient electron tunneling between the two.⁸ Aromatic compounds are typically better electron donors (*i.e.* hole receptors) due to their electron-rich π -system. With this in mind, we shall assume that the bridge/olfactant acts as an intermediate for hole transfer between a donor site and an acceptor site. Energy conservation requires that the energy transferred to vibrational motion is equal the tunneling gap.

II. THEORETICAL MODEL

For a donor-bridge-acceptor system, the tunneling matrix element for electron transfer rate can be obtained from second-order perturbation theory^{8,9}

$$V_{da}(E_{da}) = \langle A | V_{AB} (E_{da} - H_B)^{-1} V_{BD} | D \rangle \quad (1)$$

where $|A\rangle$ and $|D\rangle$ denote the final and initial states and H_B is the electronic Hamiltonian for the bridge molecule. Here, an electron can tunnel from an occupied molecular orbital on the donor, to an unoccupied molecular orbital on the bridge, and finally be transferred to an unoccupied molecular orbital on the acceptor, E_{da} is the tunneling gap and V_{ij} denotes the couplings between the donor to bridge and bridge to acceptor. One can also consider hole transfer along similar lines.

We define the couplings between the donor to bridge and bridge to acceptor as $V_{db} = J_{db}|d\rangle\langle b|$ and $V_{ba} = J_{ba}|b\rangle\langle a|$, respectively. From the Franck-Condon principle, these can be factored from the nuclear/vibrational terms. Further, let us assume that the bridge has two electronic states, denoted as $|1\rangle$ corresponding to the neutral bridge, and $|2\rangle$ corresponding to the oxidized bridge. In addition, the bridge has a set of internal molecular vibrational states local to the molecule with normal mode

frequencies, $\{\omega_q\}$. It is important to point out that the vibrational states of the neutral are not the same as those of the ion due to shift of the energy minima and (Duschinsky) rotation of the normal modes.

We can thus evaluate the transition matrix as

$$V_{da} = J_{db}J_{ba} \sum_{\mathbf{n}_2, \mathbf{n}'_1} \frac{\langle \mathbf{n}'_1 | \mathbf{n}_2 \rangle \langle \mathbf{n}_2 | \mathbf{0}_1 \rangle}{E_{da} - (\tilde{\epsilon}_2 + \hbar\omega_2 \cdot (\mathbf{n}_2 + 1/2))}, \quad (2)$$

where the sum is over all intermediate, $\mathbf{n}_2$, and final, $\mathbf{n}'_1$, vibrational states (including overtones and combinations) of the oxidized bridge molecule and we assume that the initial vibrational populations are in their ground states. The terms in the numerator are the overlap integrals between the vibrational states of the neutral and oxidized bridge. In general, these overlaps can be computed using the approach outlined in Sharp and Rosenstock^{10,11}; however, this formally involves summing over all possible combinations of initial, final, and intermediate vibrational quantum numbers including overtone modes.

To simplify this, let us consider the bridge ionization/deionization process to be purely impulsive so far as the nuclear dynamics are concerned. Upon ionization, the nuclei in B experience a sudden change in their electronic environment. Ordinarily, the vibrational wave function would begin to evolve on the Born-Oppenheimer potential, $V^{(+)}$, corresponding to this new electronic configuration. However, before nuclear motion can occur, the charge hops to the acceptor returning B to its neutral electronic state. Over the course of the charge-transfer event, the nuclei in the bridge molecule receive an impulsive “kick” due to the sudden changes in electronic state. This excitation can be imparted to the nuclear motion on the ground-state (B) potential by an instantaneous shift in the momentum of the ground-state vibrational




FIG. 2. Olfactory Discrimination of Acetophenone isomers. a) *Drosophila melanogaster* (w^{1118}) naïvely avoid ACP in a T-maze. The behavioral responses to both d5-ACP and d8-ACP are significantly stronger than to both the d3-ACP and h-ACP. ($n = 22$ groups each) (b) *Drosophila* can discriminate between d5-ACP and d3-ACP after training with electric shock. During the training session, populations of flies are exposed to one of the isotopomers paired with an electric shock. When subsequently tested in the T-maze, the flies significantly prefer the unpaired odorant over the paired odorant ($p < 0.0001$; $n = 44$ groups each). (c) *Drosophila* can discriminate between h-ACP and d3-ACP after training with electric shock.

wavefunction along the direction of $\vec{\nabla}V^{(+)}$.

$$V_{da} = \frac{J_{ab}J_{bd}}{E_{da}} \sum_{\mathbf{n}'_1} \langle \mathbf{n}'_1 | e^{i\vec{p} \cdot \vec{x}} | \mathbf{0}_1 \rangle. \quad (3)$$

It is important to recognize the $\vec{p}$ term in this expression corresponds to the momentum imparted to the nuclear motion due to the transition and that energy conservation requires $E_{da} = p^2/2m$ be the additional kinetic energy given to the internal nuclear motion of the olfactant bridge molecule. We can factor this into contributions from each normal mode and require that the momentum imparted to a given vibrational mode is proportion to the energy gradient along that direction.

The immediate advantage of this model is that we only need to consider the nuclear motion of the neutral species and in practice we can evaluate the matrix element exactly. However, it is useful to examine just the leading order terms. If the momentum kick is small, then we can expand the translation operator for each normal mode as

$$\langle \mathbf{n}'_{1q} | e^{i\vec{p} \cdot \vec{x}} | \mathbf{0}_{1q} \rangle = \delta_{0n} + i\vec{p} \cdot \langle \mathbf{n}'_{1q} | \vec{x} | \mathbf{0}_{1q} \rangle. \quad (4)$$

This first term is simply the elastic component, which is expected to dominate. The second term contains the matrix element of the position operator. This implies that to a first approximation the inelastic scattering of an electron via the bridge species excites the vibrational modes of the bridge that are also infra-red (IR) active. This is a central component of Turin's spectroscopic theory of olfaction.^{1,3,4} However, only those modes that are directed along $\vec{\nabla}V^{(+)}$ will be excited by the impulsive scattering process.

Equation (4) gives us a direct way to rapidly screen whether or not a given odorant is expected to exhibit an isotope effect using quantum chemical means. Starting from the equilibrium position of the neutral, one first

determines the vibrational normal modes and frequencies of a given olfactant. This gives the IR response of the molecule. We then determine the energy gradient of the ionized species at this geometry and project this onto the normal modes. Since E_{da} remains an unknown in our model, we simply require that this energy be distributed amongst the normal coordinates in proportion to their projection onto the energy gradient. Thus, even if a mode has a strong IR response and satisfies energy conservation, unless that mode is directly excited by the impulsive scattering process, then it will not contribute to the overall transition rate. Our approach is similar to that of Brookes *et al.*¹²; however, our approach explicitly treats the olfactant as a bridging species between the donor and acceptor and derives all relevant input from quantum chemical treatments of the bridge. This allows us to make specific predictions regarding the potential olfactory response of a given scent molecule.

As a test case, we consider acetophenone (ACP) and its deuterated isotopomers. In d3-ACP, the hydrogens on the methyl group are replaced, in d5-ACP the hydrogens on the phenyl ring are replaced, and in d8-ACP all hydrogens are replaced with deuteriums. In all cases, we first perform geometry optimizations of the neutral species followed by calculations of vibrational frequencies using the B3LYP density functional with 6-31G(d) basis set in vacuum using the NWChem quantum chemistry package.¹³ The same functional and basis set were then used to calculate the nuclear gradients and frequencies of the radical cation at the equilibrium geometry of the neutral. This approach is numerically reliable and produces accurate potentials and gradients for this system without undue effort. Moreover, this general procedure can be applied to a broad set of olfactants.

In Figure 1(b) we show the predicted IR response and

the projection of the energy gradient of the oxidized species onto the normal mode frequencies of the neutral. Briefly, the peaks around 3100 cm^{-1} correspond to the C-H stretching modes of the molecule. In the fully deuterated d8-ACP, this peak is shifted to 2300 cm^{-1} , consistent with the typical isotope shift of a C-D stretching mode. Furthermore, the IR spectra for all but the d8-ACP is void of peaks between $1800 - 3000\text{ cm}^{-1}$. Only the fully deuterated d8-ACP has any appreciable spectral density in this region. While there is little to note in comparing the intensities for the modes below 1800 cm^{-1} , the contribution to the gradient coming from the C-H and C-D stretching modes is remarkably strong. Most notable, the peak at 2300 cm^{-1} is more or less the sum of contributions from the ring and methyl C-D stretching modes.

Based upon the relative intensities of both the IR and energy gradients around 2300 cm^{-1} in Figure 1(b), the electron transfer model suggests that *D. melanogaster* should readily discriminate h-ACP and d8-ACP. In comparing the predicted IR spectra, there is almost no IR oscillator strength in the C-D stretching modes for d3- and d5-ACP while d8-ACP exhibits a fairly strong signal. However, in comparing the gradients, the d5-ACP gradient has considerably smaller projection onto the C-D stretches modes and the d3-ACP projection is almost vanishingly small. Based upon this, one predicts that there should be an ability to differentiate between h-ACP and d8-ACP. Moreover, the flies may be able to discriminate between d3-ACP and d5-ACP even though both have very weak IR signals at 2300 cm^{-1} .

III. EXPERIMENTAL TESTS

Previously, d5-ACP and d8-ACP odorants were found to be significantly more aversive to *Drosophila* than h-ACP¹. We verified this result, examining the naïve avoidance of *Drosophila w¹¹¹⁸* to 0.3% h, d3, d5, and d8 ACP using an olfactory T-Maze (Figure 2a). The avoidance response to h-ACP and d3-ACP were indistinguishable ($F_{3,80} = 11.724$, Bonferonni-Dunn *post hoc* $p = 0.276$); the responses to d5 and d8 were also not significantly different from each other ($p = 0.842$). However, the avoidance response to d5-ACP was significantly greater than the response to d3-ACP ($p = 0.0016$), suggesting deuteriums on the benzene ring provided a different odorant quality than the methyl deuteriums, leading to an increased saliency for the d5-ACP and d8-ACP. To test this hypothesis, we used negatively-reinforced olfactory learning¹⁴. Surprisingly, the *w¹¹¹⁸* flies are capable of discriminating between d5-ACP and d3-ACP after training, indicating a significant difference in perceptual quality between these odorants (Figure 2b; $t = 24.714$, $p < 0.0001$). The *w¹¹¹⁸* flies are also capable of discriminating between h-ACP and d3-ACP after training (Figure 3c; $t = 10.596$; $p < 0.001$). The conditioned avoidance in this experiment was reduced in comparison to the conditioned avoidance of d5-ACP vs. d3-ACP, in-

dicating that d3-ACP is more similar in odorant quality to h-ACP than to d5-ACP. The robust discrimination between these very closely related odorants is consistent with our model that predicts differences in the excitation of vibrational modes of the methyl and deuterium C-D bonds by the impulsive scattering.

IV. SUMMARY

It is clearly evident that *D. melanogaster* can be trained to distinguish deuterated isotopomers of various olfactants. Whether or not this is the result of a spectroscopic detection mechanism is another question. While deuteration does shift the C-H stretching mode into the otherwise empty 2300 cm^{-1} region, our model suggests that *IR response alone does not determine whether or not the flies can discriminate isotopomers*. Moreover, this effect appears to be limited to a single spectral region and does not involve the entire IR spectrum as suggested by Turin.^{3,4} A crucial test of our model would be to identify an olfactant that can not be discriminated from its deuterated isotopomers. For example, in removing an electron from ethylene, the C=C bond-length in C_2H_4^+ will be longer than in C_2H_4 (due to reduction in bond-order) but the C-H bond-lengths will be more or less unchanged. This suggests that the energy gradient in our model will be directed primarily along the C=C bond and have very little projection onto the C-H stretching modes. However, the IR spectrum of ethylene does show an isotope shift of the C-H stretching modes into the 2300 cm^{-1} region. Hence, we predict that *D. melanogaster* should *not* be able to discriminate between ethylene and its isotopomers.

Lacking detailed molecular level knowledge of the receptor sites, we speculate the tunneling gap between donor and acceptor sites is around 2300 cm^{-1} and that the narrow spectral region about E_{da} is giving rise to the observed ability for *Drosophila* to discriminate between deuterated isotopomers by scent. It is possible that the tunneling gap may be tuned to the nitrile ($\text{C}\equiv\text{N}$) stretching mode which does lie in this spectral region and may serve a role in detecting a variety of naturally occurring nitrile containing odorants.

ACKNOWLEDGMENTS

ERB was supported by the National Science Foundation (CHE-1011894) and Robert A. Welch Foundation (E-1334). The work by GR was supported by DARPA (N66001-11-1-4119).

ERB jointly conceived the study with GR, developed the theoretical model and prepared the manuscript. AC performed the quantum chemical calculations. GR and AM designed and performed the experiments.

The authors have no competing financial interests.

- ¹M. I. Franco, L. Turin, A. Merzhinov, and E. M. C. Skoulakis, "Molecular vibration-sensing component in drosophila melanogaster olfaction," *Proceedings of the National Academy of Sciences* **108**, 3797–3802 (2011), <http://www.pnas.org/content/108/9/3797.full.pdf+html>.
- ²T. Tully and W. G. Quinn, "Classical conditioning and retention in normal and mutant drosophila melanogaster." *J. Comp. Physiol. A* **157**, 263–267 (1985).
- ³L. Turin, "A spectroscopic mechanism for primary olfactory reception," *Chemical Senses* **21**, 773–791 (1996), <http://chemse.oxfordjournals.org/content/21/6/773.full.pdf+html>.
- ⁴L. TURIN, "A method for the calculation of odor character from molecular structure," *Journal of Theoretical Biology* **216**, 367 – 385 (2002).
- ⁵T. P. Hettinger, "Olfaction is a chemical sense, not a spectral sense," *Proceedings of the National Academy of Sciences* **108**, E349 (2011), <http://www.pnas.org/content/108/31/E349.full.pdf+html>.
- ⁶A. Keller and L. B. Vosshall, "A psychophysical test of the vibration theory of olfaction." *Nature Neuroscience* **7**, 337–8 (2004).
- ⁷J. Wang, Z. A. Luthe-Schulten, and K. S. Suslick, "Is the olfactory receptor a metalloprotein?" *Proceedings of the National Academy of Sciences* **100**, 3035–3039 (2003), <http://www.pnas.org/content/100/6/3035.full.pdf+html>.
- ⁸D. Devault, *Quantum mechanical tunneling in biological systems* (Cambridge University Press, Cambridge, U.K., 1984).
- ⁹O. S. Wenger, "How donor–bridge–acceptor energetics influence electron tunneling dynamics and their distance dependences," *Accounts of Chemical Research* **44**, 25–35 (2011), <http://pubs.acs.org/doi/pdf/10.1021/ar100092v>.
- ¹⁰T. E. Sharp and H. M. Rosenstock, "Franck-Condon factors for polyatomic molecules," *The Journal of Chemical Physics* **41**, 3453–3463 (1964).
- ¹¹J. WEBER and G. HOHLNEICHER, "Franck—condon factors for polyatomic molecules," *Molecular Physics* **101**, 2125–2144 (2003), <http://www.tandfonline.com/doi/pdf/10.1080/0026897031000109310>.
- ¹²J. C. Brookes, F. Hartoutsiou, A. P. Horsfield, and A. M. Stoneham, "Could humans recognize odor by phonon assisted tunneling?" *Phys. Rev. Lett.* **98**, 038101 (2007).
- ¹³M. Valiev, E. Bylaska, N. Govind, K. Kowalski, T. Straatsma, H. V. Dam, D. Wang, J. Nieplocha, E. Apra, T. Windus, and W. de Jong, "Nwchem: A comprehensive and scalable open-source solution for large scale molecular simulations," *Computer Physics Communications* **181**, 1477 – 1489 (2010).
- ¹⁴G. Roman and R. L. Davis, "Molecular biology and anatomy of drosophila olfactory associative learning," *BioEssays* **23**, 571 (2001).